

## qScript™ cDNA SuperMix

|         |           |       |                       |              |                       |
|---------|-----------|-------|-----------------------|--------------|-----------------------|
| Cat No. | 95048-025 | Size: | 25 x 20-μL reactions  | (1 x 100 μL) | <b>Store at -20°C</b> |
|         | 95048-100 |       | 100 x 20-μL reactions | (1 x 400 μL) |                       |
|         | 95048-500 |       | 500 x 20-μL reactions | (2 x 1 mL)   |                       |

### Description

qScript cDNA SuperMix provides a sensitive and easy-to-use solution for two-step RT-PCR. This 5X concentrated master mix provides all necessary components (except RNA template) for first-strand synthesis including: buffer, dNTPs, MgCl<sub>2</sub>, primers, RNase inhibitor protein, qScript reverse transcriptase and stabilizers. qScript is a RNase H(+) derivative of MMLV reverse transcriptase, optimized for reliable cDNA synthesis over a wide dynamic range of input RNA. The unique blend of oligo (dT) and random primers in the qScript cDNA SuperMix works exceptionally well with a wide variety of targets. This blend is optimized for the production of targets < 1kb in length. qScript cDNA SuperMix produces excellent results in both real-time and conventional RT-PCR.

### Components

|                       |                                                                                                                                                                                                                                     |
|-----------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| qScript cDNA SuperMix | 5X reaction buffer containing optimized concentrations of MgCl <sub>2</sub> , dNTPs (dATP, dCTP, dGTP, dTTP), recombinant RNase inhibitor protein, qScript reverse transcriptase, random primers, oligo(dT) primer and stabilizers. |
|-----------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

### Storage and Stability

qScript cDNA SuperMix is stable for 1 year when stored in a constant temperature freezer at -20°C. To extend the product's shelf-life, store the kit at -70°C. qScript cDNA SuperMix showed no loss in functional performance after 20 cycles of freezing on dry ice and thawing on ice. However, we recommend that the number of freeze-thaw cycles be kept to a minimum.

### Reaction Assembly

Place components on ice. Mix, and then briefly centrifuge to collect contents to the bottom of the tube before using.

| Component                  | Volume for 20-μL rxn. | Final Concentration       |
|----------------------------|-----------------------|---------------------------|
| qScript cDNA SuperMix (5X) | 4 μL                  | 1X                        |
| RNA template               | variable              | (1 μg to 10 pg total RNA) |
| RNase/DNase-free water     | <u>variable</u>       |                           |
| Total Volume (μL)          | 20 μL                 |                           |

**Note:** for smaller reaction volumes (i.e. 10-μL reactions), scale components proportionally.

### Reaction Protocol

- Combine reagents in 0.2-mL micro-tubes or 96-well plate sitting on ice.
- After sealing each reaction, vortex gently to mix contents. Centrifuge briefly to collect components at the bottom of the reaction tube.
- Incubate:
  - 5 minutes at 25°C
  - 30 minutes at 42°C
  - 5 minutes at 85°C
  - Hold at 4°C
- After completion of cDNA synthesis, use 1/5th to 1/10th of the first-strand reaction (2-4 μL) for PCR amplification. If desired, cDNA product can be diluted with 10 mM Tris-HCl (pH 8.0), 0.1 mM EDTA and stored at -20°C.

## Guidelines for Reverse Transcription-qPCR

**Minus RT-controls:** Accurate quantification of gene expression by RT-qPCR requires testing and reporting the extent of contamination of genomic DNA in each RNA sample for each gene of interest. The presence of trace amounts of gDNA does not usually interfere with quantification of high copy reference genes. However, it can have a significant contribution on signal for low copy genes. Even when using primers that are separated by intronic sequence or bridge exon junctions, the presence of genomic DNA can produce positive signals from amplification of pseudogene or off-target PCR product. Therefore, it is important to always include the appropriate "no RT" or "minus RT" control reactions in your experimental design.

Since the reverse transcriptase is an integral component of qScript cDNA SuperMix, it is not feasible to construct a formal cDNA synthesis control that includes all components except the RT. The most direct method to test for the presence of genomic DNA is to bypass the RT step and use an equivalent amount of the RNA preparation directly for PCR amplification. For example: if you start with 1 µg of total RNA for cDNA synthesis and use 1/10<sup>th</sup> of the first-strand reaction as template for qPCR; then use 100 ng of total RNA as template for the minus RT-control qPCR. Any signal from the RNA only reaction is attributable to the presence of genomic DNA.

**DNase digestion of total RNA:** Trace levels of genomic DNA can obscure accurate quantification, particularly when the specific gene(s) of interest are low copy. PerfeCta® DNase I is a high purity, recombinant DNase I preparation that is free of any contaminating RNases. It provides a simple and rapid solution to eliminate residual genomic DNA that is directly compatible with qScript cDNA SuperMix, or other first-strand synthesis kits. The supplied *Reaction Buffer* and proprietary *Stop Buffer* support a simple heat-kill step that permanently inactivates all trace levels of DNase activity before the cDNA synthesis step. Heat-kill procedures used by other DNase I reagents are ineffective and not compatible with qScript cDNA SuperMix. Residual, or renatured, DNase will degrade cDNA product and alter apparent expression levels. If using other sources of RNase-free DNase I, it is essential to remove all traces of DNase activity before proceeding with first-strand synthesis. Suitable RNA purification methods include phenol:chloroform extraction followed by ethanol precipitation, or the use of chaotropic salts and a silica-based RNA purification cartridge or column. Please call technical support at (800) 364-2149 or visit our web site at [www.quantabio.com](http://www.quantabio.com) if you require additional information or protocols.

## Quality Control

Kit components are free of contaminating DNase and RNase. qScript cDNA SuperMix is functionally tested in reverse transcription quantitative PCR (RT-qPCR). First-strand synthesis is performed in triplicate on each dilution of a log-fold serial dilution of HeLa cell total RNA from 1 pg to 1 µg. One-tenth of each first-strand reaction is used for qPCR amplification. Kinetic analysis must demonstrate linear resolution over five orders of dynamic range ( $r^2 > 0.995$ ) and a PCR efficiency  $> 90\%$ .

## Limited Use Label License

This product is covered by US patent 7,470,515, US patent 7,638,612 and other patents pending in the United States and Europe. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer. The buyer is not authorized to sell or otherwise transfer this product, any of its components to a third party. The purchase of this product does not authorize the purchaser to use the product or any of its components for manufacture of commercial product. For information on obtaining a license to this product for purposes other than research, contact Licensing Department, Quanta BioSciences, Inc., 202 Perry Parkway, Suite 1, Gaithersburg, MD 20877; Telephone number: 1-800-364-2149.

## Related Products

PerfeCta DNase I, Cat. No. 95150-01K

qScript is a trademark of Quanta BioSciences, Inc.

PerfeCta is a registered Trademark of Quanta BioSciences, Inc.